Recombinant Human EXOSC9 Protein (M1-N439), Tag Free

Cat. No. EXOSC9-0731H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human GG-EXOSC9(M1-N439 end) Protein was expressed in E. coli.
Species	Human
Source	E.coli
ProteinLength	M1-N439
Description	Non-catalytic component of the RNA exosome complex which has 3'->5' exoribonuclease activity and participates in a multitude of cellular RNA processing and degradation events. In the nucleus, the RNA exosome complex is involved in proper maturation of stable RNA species such as rRNA, snRNA and snoRNA, in the elimination of RNA processing by-products and non-coding 'pervasive' transcripts, such as antisense RNA species and promoter-upstream transcripts (PROMPTs), and of mRNAs with processing defects, thereby limiting or excluding their export to the cytoplasm. The RNA exosome may be involved in Ig class switch recombination (CSR) and/or Ig variable region somatic hypermutation (SHM) by targeting AICDA deamination activity to transcribed dsDNA substrates. In the cytoplasm, the RNA exosome complex is involved in general mRNA turnover and specifically degrades inherently unstable mRNAs containing AU-rich elements (AREs) within their 3' untranslated regions, and in RNA surveillance pathways, preventing translation of aberrant mRNAs. It seems to be involved in degradation of histone mRNA. The catalytic inactive RNA exosome core complex of 9 subunits (Exo-9) is proposed to play a pivotal role in the binding and presentation of RNA for ribonucleolysis, and to

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	serve as a scaffold for the association with catalytic subunits and accessory proteins or complexes. EXOSC9 binds to ARE-containing RNAs.
Form	Liquid
Endotoxin	< 0.01 EU per μg of the protein
Purity	90%
Stability	Samples are stable for up to twelve months from date of receipt at -20 to -80 centigrade.
Storage	Store it under sterile conditions at -20 to -80 centigrade. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Storage Buffer	Supplied as sterile 50 mM Tris-HCl (pH7.5), 200 mM NaCl, 20% glycerol
Shipping	It is shipped out with blue ice.

GENE INFORMATION

Gene Name	EXOSC9 exosome component 9 [Homo sapiens (human)]
Official Symbol	EXOSC9
Synonyms	EXOSC9; exosome component 9; PMSCL1, polymyositis/scleroderma autoantigen 1, 75kDa; exosome complex component RRP45; p5; p6; PM/Scl 75; polymyositis/scleroderma autoantigen 1 (75kD); RRP45; Rrp45p; autoantigen PM/Scl 1; PMSCL autoantigen, 75kD; exosome complex exonuclease RRP45; polymyositis/scleroderma autoantigen 1, 75kDa; P75 polymyositis-scleroderma overlap syndrome associated autoantigen; P75 polymyositis-scleroderma overlap

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	syndrome-associated autoantigen; PMSCL1; PM/Scl-75;
Gene ID	5393
mRNA Refseq	NM_001034194
Protein Refseq	NP_001029366
MIM	606180
UniProt ID	Q06265

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA