

Recombinant Human IL10RB Protein, Met20-Ser220, C-Fc tagged

Cat. No. IL10RB-202H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human IL-10 R beta Protein (Met20-Ser220) with Human IgG1 (Pro100-Lys330) Fc tag at C-terminus was expressed in Chinese Hamster Ovary cell line.
Species	Human
Source	CHO
ProteinLength	20-220 a.a.
Description	The protein encoded by this gene belongs to the cytokine receptor family. It is an accessory chain essential for the active interleukin 10 receptor complex. Coexpression of this and IL10RA proteins has been shown to be required for IL10-induced signal transduction. This gene and three other interferon receptor genes, IFAR2, IFNAR1, and IFNGR2, form a class II cytokine receptor gene cluster located in a small region on chromosome 21.
Form	Disulfide-linked homodimer
Bio-activity	Measured by its binding ability in a functional ELISA. Recombinant Human IL-10 R beta Fc Chimera binds Recombinant Human IL-28 R alpha/IFN-lambda R1 Fc Chimera in the presence of Recombinant Human IL-28B/IFN-lambda 3 with an ED50 of 0.250-3.00 µg/mL.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	Predicted Molecular Mass: 50 kDa SDS-PAGE: 72-82 kDa, under reducing conditions.
Endotoxin	< 0.10 EU/μg of the protein by the LAL method.
Purity	> 95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie Blue Staining.
N-terminal Sequence Analysis	Met20
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Storage Buffer	Lyophilized from a 0.2 μm filtered solution in PBS.
Shipping	The product is shipped at ambient temperature.
Reconstitution	Reconstitute at 500 μg/mL in PBS.
References	1. Commins, S. et al. (2008) J. Allergy Clin. Immunol. 121:1108. 2. Pestka, S. et al. (2004) Annu. Rev. Immunol. 22:929. 3. Kotenko, S.V. et al. (1997) EMBO J. 16:5894.

GENE INFORMATION

Gene Name	IL10RB interleukin 10 receptor, beta [Homo sapiens (human)]
Official Symbol	IL10RB

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Synonyms

IL10RB; interleukin 10 receptor, beta; CRFB4, D21S58, D21S66; interleukin-10 receptor subunit beta; CDW210B; CRF2 4; IL 10R2; IL-10RB; IL-10R subunit 2; IL-10R subunit beta; IL-10 receptor subunit beta; cytokine receptor class-II CRF2-4; interleukin-10 receptor subunit 2; cytokine receptor class-II member 4; cytokine receptor family 2 member 4; cytokine receptor family II, member 4; CRFB4; CRF2-4; D21S58; D21S66; IL-10R2

Gene ID

3588

mRNA Refseq

NM_000628

Protein Refseq


NP_000619

MIM

123889

UniProt ID

Q08334


Binding Activity


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

SDS-PAGE



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA