

Recombinant Human NLRP6 Protein (M1-F892), Tag Free

Cat. No. NLRP6-0069H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human NLRP6(M1-F892 end) Protein was expressed in Insect cell.
Species	Human
Source	Insect Cells
ProteinLength	M1-F892
Description	Acts as the sensor component of the NLRP6 inflammasome, which mediates inflammasome activation in response to various pathogen-associated signals, leading to maturation and secretion of IL1B and IL18. Inflammasomes are supramolecular complexes that assemble in the cytosol in response to pathogens and other damage-associated signals and play critical roles in innate immunity and inflammation. Acts as a recognition receptor (PRR): recognizes and binds specific pathogens and other damage-associated signals, such as lipoteichoic acid (LTA), a cell-wall component of Gram-positive bacteria, or double stranded RNA (dsRNA). May also recognize and bind lipopolysaccharide (LPS), a major component of the outer membrane of Gram-negative bacteria; however, LPS is probably not a major activator of the NLRP6 inflammasome. Following LTA- or dsRNA-binding, NLRP6 undergoes liquid-liquid phase separation (LLPS), enhancing multivalent interactions, an essential step for the formation of the NLRP6 inflammasome polymeric complex. The NLRP6 inflammasome acts by promoting recruitment of effector proinflammatory caspases (CASP1 and/or CASP4) that catalyze maturation and secretion of IL1B and IL18 in the extracellular milieu. The NLRP6 inflammasome plays a central role in the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

maintenance of epithelial integrity and host defense against microbial infections in the intestine. Required to restrict infection against Gram-positive bacteria by recognizing lipoteichoic acid (LTA), leading to recruitment of CASP4 and CASP1, and subsequent maturation and secretion of IL1B and IL18. Involved in intestinal antiviral innate immunity together with DHX15: recognizes and binds viral dsRNA to restrict infection by enteric viruses through the interferon pathway and GSDMD-dependent release of IL18. Required to prevent infection by the apicomplexan parasite *Cryptosporidium* in enterocytes by promoting GSDMD-dependent release of IL18. The NLRP6 inflammasome may also regulate the gut microbiota composition by acting as a sensor of microbiota-associated metabolites to form a PYCARD/ASC-dependent inflammasome for downstream IL18 release and secretion of antimicrobial peptides. Essential for gut mucosal self-renewal and proliferation. Regulate mucus secretion in an inflammasome- and autophagy-dependent manner to prevent invasion by enteric bacteria,. During systemic bacterial infections, the NLRP6 inflammasome negatively regulates neutrophil recruitment and neutrophil extracellular traps (NETs) formation. May promote peripheral nerve recovery following injury via an inflammasome-independent mechanism.

Form	Liquid
Endotoxin	< 0.01 EU per µg of the protein
Purity	90%
Stability	Samples are stable for up to twelve months from date of receipt at -20 to -80 centigrade.
Storage	Store it under sterile conditions at -20 to -80 centigrade. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage Buffer	Supplied as sterile 50mM Tris-HCl (pH 7.5), 200mM NaCl, 20% glycerol
Shipping	It is shipped out with blue ice.
GENE INFORMATION	
Gene Name	NLRP6 NLR family, pyrin domain containing 6 [Homo sapiens (human)]
Official Symbol	NLRP6
Synonyms	NLRP6; NLR family, pyrin domain containing 6; NACHT, leucine rich repeat and PYD containing 6, NALP6; NACHT, LRR and PYD domains-containing protein 6; CLR11.4; nucleotide binding oligomerization domain; leucine rich repeat and pyrin domain containing 6; PAN3; PYPAF5; angiotensin II/vasopressin receptor; PYRIN-containing APAF1-like protein 5; NACHT, LRR and PYD containing protein 6; NACHT, leucine rich repeat and PYD containing 6; nucleotide-binding oligomerization domain, leucine rich repeat and pyrin domain containing 6; NALP6;
Gene ID	171389
mRNA Refseq	NM_138329
Protein Refseq	NP_612202
MIM	609650
UniProt ID	P59044