

Recombinant Human SLC25A5 Protein (T2-Y297), 8×His-MBP, Flag tagged

Cat. No. SLC25A5-0077H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human 8His-MBP-TEV-(GG)-SLC25A5(T2-Y297)-Flag Protein was expressed in Insect cell.
Species	Human
Source	Insect Cells
ProteinLength	T2-Y297
Description	ADP:ATP antiporter that mediates import of ADP into the mitochondrial matrix for ATP synthesis, and export of ATP out to fuel the cell. Cycles between the cytoplasmic-open state (c-state) and the matrix-open state (m-state): operates by the alternating access mechanism with a single substrate-binding site intermittently exposed to either the cytosolic (c-state) or matrix (m-state) side of the inner mitochondrial membrane. In addition to its ADP:ATP antiporter activity, also involved in mitochondrial uncoupling and mitochondrial permeability transition pore (mPTP) activity. Plays a role in mitochondrial uncoupling by acting as a proton transporter: proton transport uncouples the proton flows via the electron transport chain and ATP synthase to reduce the efficiency of ATP production and cause mitochondrial thermogenesis. Proton transporter activity is inhibited by ADP:ATP antiporter activity, suggesting that SLC25A5/ANT2 acts as a master regulator of mitochondrial energy output by maintaining a delicate balance between ATP production (ADP:ATP antiporter activity) and thermogenesis (proton transporter activity). Proton transporter activity requires

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

free fatty acids as cofactor, but does not transport it. Probably mediates mitochondrial uncoupling in tissues that do not express UCP1. Also plays a key role in mPTP opening, a non-specific pore that enables free passage of the mitochondrial membranes to solutes of up to 1.5 kDa, and which contributes to cell death. It is however unclear if SLC25A5/ANT2 constitutes a pore-forming component of mPTP or regulates it. Acts as a regulator of mitophagy independently of ADP:ATP antiporter activity: promotes mitophagy via interaction with TIMM44, leading to inhibit the presequence translocase TIMM23, thereby promoting stabilization of PINK1. As part of the mitotic spindle-associated MMXD complex it may play a role in chromosome segregation.

Form	Liquid
Endotoxin	< 0.01 EU per µg of the protein
Purity	80%
Stability	Samples are stable for up to twelve months from date of receipt at -20 to -80 centigrade.
Storage	Store it under sterile conditions at -20 to -80 centigrade. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Shipping	It is shipped out with blue ice.

GENE INFORMATION

Gene Name	SLC25A5 solute carrier family 25 (mitochondrial carrier; adenine nucleotide translocator), member 5 [Homo sapiens (human)]
Official Symbol	SLC25A5

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Synonyms	SLC25A5; solute carrier family 25 (mitochondrial carrier; adenine nucleotide translocator), member 5; ANT2; ADP/ATP translocase 2; 2F1; T2; T3; ANT 2; ADP/ATP carrier protein; ADP,ATP carrier protein 2; solute carrier family 25 member 5; ADP,ATP carrier protein, fibroblast isoform; adenine nucleotide translocator 2 (fibroblast); AAC2;
Gene ID	292
mRNA Refseq	NM_001152
Protein Refseq	NP_001143
MIM	300150
UniProt ID	P05141