

Active Recombinant Human TNFRSF4, HIgG1 Fc-tagged

Cat. No. TNFRSF4-521H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	The extracellular domain of human CD134 (NP_003318.1) (Leu29-Ala216) is fused to the N-terminus of the Fc region of human IgG1 was expressed in CHO cell.
Species	Human
Source	CHO
ProteinLength	29-216 a.a.
Description	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor has been shown to activate NF-kappaB through its interaction with adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that this receptor promotes the expression of apoptosis inhibitors BCL2 and BCL2L1/BCL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation.
Form	Lyophilized from 0.2µm-filtered solution in PBS.
Bio-activity	Measured by its binding ability in a functional ELISA.
Molecular Mass	46KDa (monomer)
AA Sequence	Leu29-Ala216

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin	<0.06 eu/μg="" as="" determined="" by="" lal="" test="">
Purity	>98%, by SDS-PAGE under reducing conditions.
Stability	Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.
Reconstitution	Reconstitute at 100μg/ml in sterile PBS.
Warning	Avoid freeze/thaw cycles.

GENE INFORMATION

Gene Name	TNFRSF4 tumor necrosis factor receptor superfamily, member 4 [Homo sapiens]
Official Symbol	TNFRSF4
Synonyms	TNFRSF4; tumor necrosis factor receptor superfamily, member 4; TXGP1L; tumor necrosis factor receptor superfamily member 4; ACT35; CD134; OX40; OX40 antigen; ACT35 antigen; ATC35 antigen; CD134 antigen; OX40 homologue; OX40L receptor; OX40 cell surface antigen; lymphoid activation antigen ACT35; TAX transcriptionally-activated glycoprotein 1 receptor; tax-transcriptionally activated glycoprotein 1 receptor;
Gene ID	7293
mRNA Refseq	NM_003327
Protein Refseq	NP_003318
MIM	600315

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	P43489
Chromosome Location	1p36
Pathway	Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Downstream signaling in naive CD8+ T cells, organism-specific biosystem;
Function	binding; receptor activity; tumor necrosis factor-activated receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA