

Recombinant Human UBE2D3 Protein (A2-M147), Tag Free

Cat. No. UBE2D3-0031H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human (GG)-UB2D3(A2-M147 end) Protein was expressed in Insect cell.
Species	Human
Source	Insect Cells
ProteinLength	A2-M147
Description	Accepts ubiquitin from the E1 complex and catalyzes its covalent attachment to other proteins. In vitro catalyzes 'Lys-11', as well as 'Lys-48'-linked polyubiquitination. Cooperates with the E2 CDC34 and the SCF(FBXW11) E3 ligase complex for the polyubiquitination of NFKBIA leading to its subsequent proteasomal degradation. Acts as an initiator E2, priming the phosphorylated NFKBIA target at positions 'Lys-21' and/or 'Lys-22' with a monoubiquitin. Ubiquitin chain elongation is then performed by CDC34, building ubiquitin chains from the UBE2D3-primed NFKBIA-linked ubiquitin. Acts also as an initiator E2, in conjunction with RNF8, for the priming of PCNA. Monoubiquitination of PCNA, and its subsequent polyubiquitination, are essential events in the operation of the DNA damage tolerance (DDT) pathway that is activated after DNA damage caused by UV or chemical agents during S-phase. Associates with the BRCA1/BARD1 E3 ligase complex to perform ubiquitination at DNA damage sites following ionizing radiation leading to DNA repair. Targets DAPK3 for ubiquitination which influences promyelocytic leukemia protein nuclear body (PML-NB) formation in the nucleus. In conjunction with the MDM2 and TOPORS E3 ligases, functions

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

ubiquitination of p53/TP53. Supports NRDP1-mediated ubiquitination and degradation of ERBB3 and of BRUCE which triggers apoptosis. In conjunction with the CBL E3 ligase, targets EGFR for polyubiquitination at the plasma membrane as well as during its internalization and transport on endosomes. In conjunction with the STUB1 E3 quality control E3 ligase, ubiquitinates unfolded proteins to catalyze their immediate destruction. Together with RNF135, catalyzes the viral RNA-dependent 'Lys-63'-linked polyubiquitination of RIG-I/DDX58 to activate the downstream signaling pathway that leads to interferon beta production.

Form	Liquid
Endotoxin	< 0.01 EU per µg of the protein
Purity	85%
Stability	Samples are stable for up to twelve months from date of receipt at -20 to -80 centigrade.
Storage	Store it under sterile conditions at -20 to -80 centigrade. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Storage Buffer	Supplied as sterile 50 mM Tris-HCl (pH7.5), 200 mM NaCl, 5% glycerol, 1 mM DTT.
Shipping	It is shipped out with blue ice.

GENE INFORMATION

Gene Name	UBE2D3 ubiquitin-conjugating enzyme E2D 3 [Homo sapiens (human)]
Official Symbol	UBE2D3

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Synonyms	UBE2D3; ubiquitin-conjugating enzyme E2D 3; ubiquitin conjugating enzyme E2D 3 (homologous to yeast UBC4/5), ubiquitin conjugating enzyme E2D 3 (UBC4/5 homolog, yeast); ubiquitin-conjugating enzyme E2 D3; Ubch5C; E2(17)KB 3; ubiquitin-protein ligase D3; ubiquitin carrier protein D3; ubiquitin-conjugating enzyme E2(17)KB 3; ubiquitin-conjugating enzyme E2-17 kDa 3; ubiquitin-conjugating enzyme E2D 3 (UBC4/5 homolog, yeast); ubiquitin-conjugating enzyme E2D 3 (homologous to yeast UBC4/5); UBC4/5; UBCH5C; E2(17)KB3; MGC5416; MGC43926;
Gene ID	7323
mRNA Refseq	NM_003340
Protein Refseq	NP_003331
MIM	602963
UniProt ID	P61077