Recombinant Tropaeolum majus O-acyltransferase Protein, N-10×His tagged

Cat. No. O-acyltransferase-07T **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Tropaeolum majus O-acyltransferase Protein with N-10×His tag was expressed in E. coli.
Species	Tropaeolum majus
Source	E.coli
Molecular Mass	60.3 kDa
AA Sequence	MAVAESSQNTTTMSGHGDSDLNNFRRRKPSSSVIEPSSSGFTSTNGVPATGHVAE NRDQDRVGAMENATGSVNLIGNGGGVVIGNEEKQVGETDIRFTYRPSFPAHRRVRE SPLSSDAIFKQSHAGLFNLCIVVLIAVNSRLIIENLMKYGWLIDTGFWFSSRSLGDWSI FMCCLTLPIFPLAAFIVEKLVQRNHISELVAVLLHVIVSTAAVLYPVIVILTCDSVYMSGV VLMLFGCIMWLKLVSYAHTSSDIRTLAKSGYKGDAHPNSTIVSCSYDVSLKSLAYFM VAPTLCYQPSYPRSSCIRKGWVVRQFVKLIVFIGLMGFIIEQYINPIVRNSKHPLKGDF LYAIERVLKLSVPNLYVWLCMFYSFFHLWLNILAELLRFGDREFYKDWWNAKTVAEY WKMWNMPVHRWMVRHLYFPCLRNGIPKEGAIIIAFLVSGAFHELCIAVPCHVFKLWA FIGIMFQVPLVLITNYLQEKFSNSMVGNMIFWFIFCILGQPMCVLLYYHDLINLKEK
Purity	≥90% by SDS-PAGE
Applications	MAVAESSQNTTTMSGHGDSDLNNFRRRKPSSSVIEPSSSGFTSTNGVPATGHVAE NRDQDRVGAMENATGSVNLIGNGGGVVIGNEEKQVGETDIRFTYRPSFPAHRRVRE SPLSSDAIFKQSHAGLFNLCIVVLIAVNSRLIIENLMKYGWLIDTGFWFSSRSLGDWSI

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	FMCCLTLPIFPLAAFIVEKLVQRNHISELVAVLLHVIVSTAAVLYPVIVILTCDSVYMSGV VLMLFGCIMWLKLVSYAHTSSDIRTLAKSGYKGDAHPNSTIVSCSYDVSLKSLAYFM VAPTLCYQPSYPRSSCIRKGWVVRQFVKLIVFIGLMGFIIEQYINPIVRNSKHPLKGDF LYAIERVLKLSVPNLYVWLCMFYSFFHLWLNILAELLRFGDREFYKDWWNAKTVAEY WKMWNMPVHRWMVRHLYFPCLRNGIPKEGAIIIAFLVSGAFHELCIAVPCHVFKLWA FIGIMFQVPLVLITNYLQEKFSNSMVGNMIFWFIFCILGQPMCVLLYYHDLINLKEK
Storage	Store it under sterile conditions at -20 to -80 centigrade. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Storage Buffer	Lyophilized from PBS,0.05% Brij-78, 6% Trehalose, pH 7.4 The volume before lyophilization is 100ul/vial.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20/-80 centigrade. Our default final concentration of glycerol is 50%. Customers could use it as reference.

GENE INFORMATION

Official Symbol	O-acyltransferase
Synonyms	O-acyltransferase; OATs

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

SDS-PAGE	 Observed band size: Monomer: 60 kDa It is speculated that the protein forms a dimeric structure. (Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel. Boiling SDS-treated samples may result in membrane protein aggregation. Incubation at 37 ° C for 10 min is an effective alternative.

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA