

Active Recombinant Human ICAM1 Protein (28-480aa), C-hlgG-His-tagged

Cat. No. ICAM1-14H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant human ICAM-1/CD54 (28-480aa), fused to hlgG-His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.
Species	Human
Source	Insect Cells
ProteinLength	28-480 aa
Description	ICAM-1/CD54, also known as intercellular adhesion molecule 1, is a member of the immunoglobulin superfamily. Its expression is weak on leukocytes, epithelial and resting endothelial cells, as well as some other cell types, but expression can be stimulated by IFNG, TNFa, IL-1b and LPS. They are important in inflammation, immune responses and in intracellular signalling events. It is known to bind to leucocyte integrins CD11/CD18 such as LFA-1 and Macrophage-1 antigen, during inflammation and in immune responses. It has been implicated in subarachnoid hemorrhage and expressed by respiratory epithelial cells is also the binding site for rhinovirus, the causative agent of most common colds.
Form	Liquid
Bio-activity	Measured by the ability of the immobilized protein to support the adhesion of HL-60 human promyelocytic cells. When cells are added to Human ICAM-1/CD54 coated

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


	plates, the ED50 range $\leq 2 \mu\text{g/mL}$.
Molecular Mass	76.5 kDa (692 aa)
AA Sequence	QTSVSPSKVILPRGGSVLVTCSTSCDQPKLLGIETPLPKKELLPGNNRKVYELSNVQ EDSQPMCYSNCPDGQSTAKTFLTVYWTPERVELAPLPSWQPVGKNLTLRCQVEGG APRANLTVLLRGEKELKREPAVGEPAEVTTTVLVRRDHHGANFSCRTDLRPQG LLEFENTSAPYQLQTFVLPATPPQLVSPRVLEVDQTQGTVCSLDGLFPVSEAQVHLA LGDQRLNPTVTYGNDSFSAKASVSVTAEDEGTQRLTCAVILGNQSQETLQTVTIYSF PAPNVILTKPEVSEGTEVTVKCEAHPRAKVTLNGVPAQPLGPRAQLLLKATPEDNGR SFSCSATLEVAGQLIHKNQTRRLRVLYGPRLDERDCPGNWTWPENSQQTPMCQA WGNPLPELKCLKDGTFLPIGESVTVTRDLEGTYLCRARSTQGEVTREVTNVLSPR YE
Endotoxin	< 1 EU/ μg of protein (determined by LAL method)
Purity	> 90% by SDS-PAGE
Applications	SDS-PAGE, Bioactivity
Notes	For research use only. This product is not intended or approved for human, diagnostics or veterinary use.
Storage	Can be stored at +2 to +8 centigrade for 1 week. For long term storage, aliquot and store at -20 to -80 centigrade. Avoid repeated freezing and thawing cycles.
Concentration	0.25 mg/mL (determined by Bradford assay)
Storage Buffer	Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

GENE INFORMATION

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene Name	ICAM1 intercellular adhesion molecule 1 [Homo sapiens (human)]
Official Symbol	ICAM1
Synonyms	ICAM1; intercellular adhesion molecule 1; BB2; CD54; human rhinovirus receptor; ICAM-1; cell surface glycoprotein P3.58; major group rhinovirus receptor; intercellular adhesion molecule 1 (CD54), human rhinovirus receptor; P3.58
Gene ID	3383
mRNA Refseq	NM_000201
Protein Refseq	NP_000192
MIM	147840
UniProt ID	P05362
Biological Activity	 Human ICAM-1/CD54 stimulates cell adhesion of the HL-60 human promyelocytic

cells. The ED50 range $\leq 2 \mu\text{g/mL}$.

SDS-PAGE



3 μg by SDS-PAGE under reducing condition and visualized by coomassie blue stain

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA