

Active Recombinant Mouse Icosl, Fc-tagged, mutant

Cat. No. Icosl-356M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	The extracellular domain of mouse CD275 [B7-H2] (NP_056605.1)(Thr 48-Lys279) is fused to the N-terminus of the Fc region of a mutant mouse IgG2a.
Species	Mouse
Source	CHO
ProteinLength	48-279 a.a.
Form	Lyophilized from 0.2µm-filtered solution in PBS.
Bio-activity	Measured by its binding ability in a functional ELISA. Immobilized human ICOS at 1 µg/ml (100µl/well) can bind biotinylated B7-H2(mouse):Fc (mouse) (rec.) (non-lytic) with a linear range of 0.1 - 1.0 µg/ml. Optimal dilutions should be determined by each I
Endotoxin	<0.06 eu/µg="" as="" determined="" by="" lal="">
Purity	≥98%, by SDS-PAGE under reducing conditions.
Stability	Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.
Reconstitution	Reconstitute at 100µg/ml in sterile PBS.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Warning Avoid freeze/thaw cycles.

GENE INFORMATION

Gene Name [Icosl](#)

Official Symbol Icosl

Synonyms B7h; GI50; GL50; B7-H2; LICOS; B7RP-1; GL50-B; ICOS-L; Icoslg; Ly115l; AU044799; BG071784; mKIAA0653

Gene ID [50723](#)

mRNA Refseq [NM_015790](#)

Protein Refseq [NP_056605](#)

UniProt ID [Q9JHJ8](#)

Chromosome Location 10 C1; 10

Pathway Adaptive Immune System, organism-specific biosystem; Cell adhesion molecules (CAMs), conserved biosystem

Function protein binding, receptor binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA