

Recombinant Mouse FGFR2 cell lysate

Cat. No. FGFR2-001MCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Mouse FGFR2 / CD332 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer.
Species	Mouse
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube.2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min.3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles.Notes:The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	Fgfr2 fibroblast growth factor receptor 2 [Mus musculus]
Official Symbol	FGFR2
Synonyms	FGFR2; fibroblast growth factor receptor 2; keratinocyte growth factor receptor; Bek; svs; KGFR; Fgfr7; Fgfr-2; Fgfr-7; KGFRTr; AU043015; AW556123; MGC102519;
Gene ID	14183
mRNA Refseq	NM_010207
Protein Refseq	NP_034337
Pathway	Downstream signaling of activated FGFR, organism-specific biosystem; ESC Pluripotency Pathways, organism-specific biosystem; Endocytosis, organism-specific biosystem; Endocytosis, conserved biosystem; FGFR ligand binding and activation, organism-specific biosystem; FGFR2 ligand binding and activation, organism-specific biosystem; FGFR2b ligand binding and activation, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Function

ATP binding; fibroblast growth factor 1 binding; fibroblast growth factor binding; fibroblast growth factor binding; fibroblast growth factor-activated receptor activity; heparin binding; kinase activity; nucleotide binding; protein binding; protein homodimerization activity; protein kinase activity; protein tyrosine kinase activity; receptor activity; transferase activity; transferase activity, transferring phosphorus-containing groups; transmembrane receptor protein tyrosine kinase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA