

Recombinant Human CGA cell lysate

Cat. No. CGA-001HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human FSH alpha / TSH alpha / CGA derived in Human cells. The whole cell lysate is provided in 1X Sample Buffer.
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube.2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min.3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles.Notes:The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	CGA glycoprotein hormones, alpha polypeptide [Homo sapiens]
Official Symbol	CGA
Synonyms	CGA; glycoprotein hormones, alpha polypeptide; glycoprotein hormones alpha chain; chorionic gonadotropin; alpha polypeptide; follicle stimulating hormone alpha subunit; FSHA; GPHA; GPHA1; HCG; LHA; luteinizing hormone alpha chain; lutropin alpha chain; thyroid stimulating hormone alpha chain; TSHA; FSH-alpha; LSH-alpha; TSH-alpha; follitropin alpha chain; thyrotropin alpha chain; choriogonadotropin alpha chain; chorionic gonadotrophin subunit alpha; thyroid-stimulating hormone alpha chain; follicle-stimulating hormone alpha chain; chorionic gonadotropin, alpha polypeptide; follicle-stimulating hormone alpha subunit; anterior pituitary glycoprotein hormones common subunit alpha; CG-ALPHA;
Gene ID	1081
mRNA Refseq	NM_000735

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq	NP_000726
MIM	118850
UniProt ID	P01215
Chromosome Location	6q14-q21
Pathway	Amine-derived hormones, organism-specific biosystem; Androgen biosynthesis, organism-specific biosystem; Autoimmune thyroid disease, organism-specific biosystem; Autoimmune thyroid disease, conserved biosystem; Class A/1 (Rhodopsin-like receptors), organism-specific biosystem; G alpha (s) signalling events, organism-specific biosystem; GPCR downstream signaling, organism-specific biosystem;
Function	hormone activity; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA