

Recombinant Cynomolgus TNFSF11 cell lysate

Cat. No. TNFSF11-001CCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Cynomolgus RANKL / OPGL / TNFSF11 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer.
Species	Cynomolgus
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube.2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min.3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles.Notes:The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.
GENE INFORMATION	
Gene Name	TNFSF11 tumor necrosis factor (ligand) superfamily, member 11 [<i>Macaca fascicularis</i>]
Official Symbol	TNFSF11
Synonyms	tumor necrosis factor (ligand) superfamily, member 11; Receptor activator of nuclear factor kappa-B ligand
Gene ID	102137946
mRNA Refseq	XM_005585744
Protein Refseq	XP_005585801
Chromosome Location	chromosome: 17
Pathway	Cytokine-cytokine receptor interaction, organism-specific biosystem; Prolactin

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

signaling pathway, organism-specific biosystem; Rheumatoid arthritis, conserved biosystem

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA