

Recombinant Human NCF2 cell lysate

Cat. No. NCF2-001HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human NCF2 / NCF-2 / P67phox derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.
GENE INFORMATION	
Gene Name	NCF2 neutrophil cytosolic factor 2 [Homo sapiens]
Official Symbol	NCF2
Synonyms	NCF2; neutrophil cytosolic factor 2; neutrophil cytosolic factor 2 (65kD, chronic granulomatous disease, autosomal 2); neutrophil cytosol factor 2; chronic granulomatous disease; autosomal 2; NADPH oxidase activator 2; NOXA2; p67phox; 67 kDa neutrophil oxidase factor; neutrophil NADPH oxidase factor 2; chronic granulomatous disease, autosomal 2; NCF-2; P67PHOX; P67-PHOX; FLJ93058;
Gene ID	4688
mRNA Refseq	NM_000433
Protein Refseq	NP_000424
MIM	608515

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	P19878
Chromosome Location	1q25
Pathway	Adaptive Immune System, organism-specific biosystem; Antigen processing-Cross presentation, organism-specific biosystem; Class I MHC mediated antigen processing and presentation, organism-specific biosystem; Cross-presentation of particulate exogenous antigens (phagosomes), organism-specific biosystem; Immune System, organism-specific biosystem; Leishmaniasis, organism-specific biosystem;
Function	Rac GTPase binding; binding; electron carrier activity; protein C-terminus binding; protein binding; superoxide-generating NADPH oxidase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA