

Recombinant Mouse CDK1 cell lysate

Cat. No. CDK1-001MCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Mouse CDC2 Kinase / CDK1 derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Mouse
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

[Cdk1 cyclin-dependent kinase 1 \[Mus musculus \]](#)

Official Symbol

CDK1

Synonyms

CDK1; cyclin-dependent kinase 1; p34 protein kinase; cell division protein kinase 1; cell division cycle 2 homolog A; cell cycle p34 CDC2 kinase protein; cell division cycle control protein 2a; cell division control protein 2 homolog; Cdc2; Cdc2a; p34
cdc2="">

Gene ID

[12534](#)

mRNA Refseq

[NM_007659](#)

Protein Refseq

[NP_031685](#)

Pathway

APC/C-mediated degradation of cell cycle proteins, organism-specific biosystem;
APC/C:Cdc20 mediated degradation of Cyclin B, organism-specific biosystem;
APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

ARMS-mediated activation, organism-specific biosystem; Activated TLR4 signalling, organism-specific biosystem; Activation of APC/C and APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem; Androgen Receptor Signaling Pathway, organism-specific biosystem;

Function

ATP binding; Hsp70 protein binding; RNA polymerase II carboxy-terminal domain kinase activity; cyclin binding; cyclin-dependent protein kinase activity; histone kinase activity; kinase activity; nucleotide binding; protein binding; protein kinase activity; protein serine/threonine kinase activity; transferase activity; transferase activity, transferring phosphorus-containing groups;