

Recombinant Mouse MAP2K1 cell lysate

Cat. No. MAP2K1-001MCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Mouse MAP2K1 / MEK1 / MAPKK1 derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Mouse
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.
GENE INFORMATION	
Gene Name	Map2k1 mitogen-activated protein kinase kinase 1 [Mus musculus]
Official Symbol	MAP2K1
Synonyms	MAP2K1; mitogen-activated protein kinase kinase 1; dual specificity mitogen-activated protein kinase kinase 1; MEK 1; MAPKK 1; MAPK/ERK kinase 1; MAP kinase kinase 1; ERK activator kinase 1; mitogen activated protein kinase kinase 1; protein kinase, mitogen activated, kinase 1, p45; Mek1; MEKK1; MAPKK1; Prkmk1;
Gene ID	26395
mRNA Refseq	NM_008927
Protein Refseq	NP_032953
Pathway	ARMS-mediated activation, organism-specific biosystem; Activated TLR4 signalling, organism-specific biosystem; Acute myeloid leukemia, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Acute myeloid leukemia, conserved biosystem; Androgen Receptor Signaling Pathway, organism-specific biosystem; Axon guidance, organism-specific biosystem; B Cell Receptor Signaling Pathway, organism-specific biosystem;

Function

ATP binding; MAP kinase kinase activity; MAP kinase kinase activity; Ras GTPase binding; kinase activity; mitogen-activated protein kinase kinase kinase binding; nucleotide binding; protein kinase activity; protein kinase binding; protein serine/threonine kinase activity; protein tyrosine kinase activity; receptor signaling protein tyrosine phosphatase activity; transferase activity; transferase activity, transferring phosphorus-containing groups;