

Recombinant Human SAE1 cell lysate

Cat. No. SAE1-001HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human SAE1 derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	SAE1 SUMO1 activating enzyme subunit 1 [Homo sapiens]
Official Symbol	SAE1
Synonyms	SAE1; SUMO1 activating enzyme subunit 1; SUMO-activating enzyme subunit 1; activator Of sumo 1; AOS1; FLJ3091; Sua1; activator of SUMO1; SUMO-1 activating enzyme subunit 1; sentrin/SUMO-activating protein AOS1; SUMO-1 activating enzyme E1 N subunit; ubiquitin-like 1-activating enzyme E1A; ubiquitin-like protein SUMO-1 activating enzyme; SUA1; UBLE1A; HSPC140;
Gene ID	10055
mRNA Refseq	NM_001145713
Protein Refseq	NP_001139185
MIM	613294

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	Q9UBE0
Chromosome Location	19q13.32
Pathway	Adaptive Immune System, organism-specific biosystem; Antigen processing: Ubiquitination and Proteasome degradation, organism-specific biosystem; Class I MHC mediated antigen processing & presentation, organism-specific biosystem; Immune System, organism-specific biosystem; Ubiquitin mediated proteolysis, organism-specific biosystem;
Function	ATP-dependent protein binding; contributes_to SUMO activating enzyme activity; enzyme activator activity; ligase activity; nucleotide binding; protein C-terminus binding; protein binding; protein heterodimerization activity; protein heterodimerization activity; ubiquitin activating enzyme activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA