

Recombinant Human PTN cell lysate

Cat. No. PTN-001HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human Pleiotrophin / PTN derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	PTN pleiotrophin [Homo sapiens]
Official Symbol	PTN
Synonyms	PTN; pleiotrophin; NEGF1, neurite growth promoting factor 1; HBGF8; HBNF; heparin binding growth factor 8; HBBM; OSF-1; HB-GAM; HBGF-8; HBNF-1; osteoblast-specific factor 1; heparin-binding brain mitogen; heparin-binding growth factor 8; heparin affin regulatory protein; neurite growth-promoting factor 1; heparin-binding growth-associated molecule; heparin-binding neurite outgrowth-promoting factor 1; pleiotrophin (heparin binding growth factor 8, neurite growth-promoting factor 1); HARP; NEGF1;
Gene ID	5764
mRNA Refseq	NM_002825
Protein Refseq	NP_002816

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



MIM	162095
UniProt ID	P21246
Chromosome Location	7q33
Pathway	Syndecan-3-mediated signaling events, organism-specific biosystem;
Function	growth factor activity; heparin binding; protein phosphatase inhibitor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA