

Recombinant Human KDM1A cell lysate

Cat. No. KDM1A-001HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human KDM1 / LSD1 derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	KDM1A lysine (K)-specific demethylase 1A [Homo sapiens]
Official Symbol	KDM1A
Synonyms	KDM1A; lysine (K)-specific demethylase 1A; amine oxidase (flavin containing) domain 2 , AOF2, KDM1, lysine (K) specific demethylase 1; lysine-specific histone demethylase 1A; BHC110; KIAA0601; LSD1; lysine (K)-specific demethylase 1; BRAF35-HDAC complex protein BHC110; lysine-specific histone demethylase 1; amine oxidase (flavin containing) domain 2; FAD-binding protein BRAF35-HDAC complex, 110 kDa subunit; flavin-containing amine oxidase domain-containing protein 2; AOF2; KDM1;
Gene ID	23028
mRNA Refseq	NM_001009999
Protein Refseq	NP_001009999

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	609132
UniProt ID	O60341
Chromosome Location	1p36.12
Pathway	Coregulation of Androgen receptor activity, organism-specific biosystem; Factors involved in megakaryocyte development and platelet production, organism-specific biosystem; Hemostasis, organism-specific biosystem; Notch signaling pathway, organism-specific biosystem; Notch-mediated HES/HEY network, organism-specific biosystem;
Function	MRF binding; androgen receptor binding; chromatin binding; demethylase activity; enzyme binding; flavin adenine dinucleotide binding; histone demethylase activity; histone demethylase activity (H3-K4 specific); histone demethylase activity (H3-K9 specific); histone demethylase activity (H3-dimethyl-K4 specific); ligand-dependent nuclear receptor transcription coactivator activity; oxidoreductase activity; p53 binding; protein binding; transcription factor binding; transcription regulatory region DNA binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA