

Recombinant Human PDGFRA cell lysate

Cat. No. PDGFRA-001HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human PDGFRA / CD140a derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	PDGFRA platelet-derived growth factor receptor, alpha polypeptide [Homo sapiens]
Official Symbol	PDGFRA
Synonyms	PDGFRA; platelet-derived growth factor receptor, alpha polypeptide; platelet-derived growth factor receptor alpha; CD140a; PDGFR2; PDGFR-alpha; PDGF-R-alpha; CD140a antigen; PDGFRA/BCR fusion; CD140 antigen-like family member A; platelet-derived growth factor receptor 2; alpha-type platelet-derived growth factor receptor; rearranged-in-hypereosinophilia-platelet derived growth factor receptor alpha fusion protein; CD140A; PDGFR-2; RHEPDGFRA; MGC74795;
Gene ID	5156
mRNA Refseq	NM_006206
Protein Refseq	NP_006197
MIM	173490

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	P16234
Chromosome Location	4q12
Pathway	ATF-2 transcription factor network, organism-specific biosystem; Calcium signaling pathway, organism-specific biosystem; Calcium signaling pathway, conserved biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Downstream signal transduction, organism-specific biosystem; Endocytosis, organism-specific biosystem;
Function	ATP binding; nucleotide binding; phosphatidylinositol 3-kinase binding; platelet-derived growth factor alpha-receptor activity; platelet-derived growth factor alpha-receptor activity; platelet-derived growth factor binding; platelet-derived growth factor binding; platelet-derived growth factor receptor binding; protein homodimerization activity; protein tyrosine kinase activity; receptor activity; transmembrane receptor protein tyrosine kinase activity; vascular endothelial growth factor binding; vascular endothelial growth factor-activated receptor activity;