

Recombinant Human PPP3CA cell lysate

Cat. No. PPP3CA-001HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human PPP3CA / derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

PPP3CA protein phosphatase 3, catalytic subunit, alpha isozyme [Homo sapiens]

Official Symbol

PPP3CA

Synonyms

PPP3CA; protein phosphatase 3, catalytic subunit, alpha isozyme; CALN, CALNA, protein phosphatase 3 (formerly 2B), catalytic subunit, alpha isoform , protein phosphatase 3 (formerly 2B), catalytic subunit, alpha isoform (calcineurin A alpha); serine/threonine-protein phosphatase 2B catalytic subunit alpha isoform; calcineurin A alpha; CNA1; PPP2B; protein phosphatase 2B; catalytic subunit; alpha isoform; CAM-PRP catalytic subunit; calmodulin-dependent calcineurin A subunit alpha isoform; protein phosphatase 2B, catalytic subunit, alpha isoform; protein phosphatase 3 (formerly 2B), catalytic subunit, alpha isoform; CALN; CCN1; CALNA; CALNA1;

Gene ID

5530

mRNA Refseq

NM_000944

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq	NP_000935
MIM	114105
UniProt ID	Q08209
Chromosome Location	4q24
Pathway	Alzheimers disease, organism-specific biosystem; Alzheimers disease, conserved biosystem; Amphetamine addiction, organism-specific biosystem; Amphetamine addiction, conserved biosystem; Amyotrophic lateral sclerosis (ALS), organism-specific biosystem; Amyotrophic lateral sclerosis (ALS), conserved biosystem; Apoptosis, organism-specific biosystem;
Function	calcium ion binding; calmodulin binding; calmodulin binding; calmodulin-dependent protein phosphatase activity; drug binding; enzyme binding; hydrolase activity; protein binding; protein dimerization activity; protein heterodimerization activity; protein serine/threonine phosphatase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA