

Recombinant Human EPHA4 cell lysate

Cat. No. EPHA4-001HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human EPHA4 / HEK8 (aa 570-986) derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
ProteinLength	570-986 a.a.
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

EPHA4 EPH receptor A4 [Homo sapiens]

Official Symbol

EPHA4

Synonyms

EPHA4; EPH receptor A4; EphA4 , TYRO1; ephrin type-A receptor 4; Hek8; EK8; EPH-like kinase 8; TYRO1 protein tyrosine kinase; tyrosine-protein kinase TYRO1; tyrosine-protein kinase receptor SEK; receptor protein-tyrosine kinase HEK8; SEK; HEK8; TYRO1;

Gene ID

2043

mRNA Refseq

NM_004438

Protein Refseq

NP_004429

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	602188
UniProt ID	P54764
Chromosome Location	2q36.3
Pathway	Axon guidance, organism-specific biosystem; Axon guidance, conserved biosystem; EPHA forward signaling, organism-specific biosystem; EphrinA-EPHA pathway, organism-specific biosystem;
Function	ATP binding; DH domain binding; GPI-linked ephrin receptor activity; PH domain binding; ephrin receptor binding; nucleotide binding; protein binding; protein kinase activity; receptor activity; transmembrane receptor protein tyrosine kinase activity; transmembrane-ephrin receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA