Recombinant Mouse DDR1 cell lysate

Cat. No. DDR1-401MCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Mouse DDR1 Kinase / MCK10 / CD167 derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer.Browse all transfected cell lysate positive controls
Species	Mouse
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube.2. Re-dissolve the pellet using 200μL pure water and boiled for 2-5 min.3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles.Notes:The lysate is ready to load on

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	Ddr1 discoidin domain receptor family, member 1 [Mus musculus]
Official Symbol	DDR1
Synonyms	DDR1; discoidin domain receptor family, member 1; epithelial discoidin domain-containing receptor 1; tyrosine kinase DDR; cell adhesion kinase; tyrosine-protein kinase CAK; protein-tyrosine kinase MPK-6; CD167 antigen-like family member A; discoidin receptor tyrosine kinase; epithelial discoidin domain receptor 1; Cak; Nep; PTK3A; CD167a; AI323681; 6030432F18;
Gene ID	12305
mRNA Refseq	NM_001198831
Protein Refseq	NP_001185760
Function	ATP binding; collagen binding; kinase activity; nucleotide binding; protein kinase

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	activity; protein tyrosine kinase activity; protein tyrosine kinase collagen receptor activity; receptor activity; transferase activity; transferase activity, transferring phosphorus-containing groups; transmembrane receptor protein tyrosine kinase activity;

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA