

Recombinant Human TP63 cell lysate

Cat. No. TP63-472HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human P63 / TP63 / Tumor derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

TP63 tumor protein p63 [Homo sapiens]

Official Symbol

TP63

Synonyms

TP63; tumor protein p63; TP53CP, TP53L, TP73L, tumor protein p53 competing protein , tumor protein p53 like , tumor protein p73 like; tumor protein 63; EEC3; KET; NBP; OFC8; p51; p53CP; p63; p73H; p73L; SHFM4; CUSP; transformation-related protein 63; tumor protein p53-competing protein; amplified in squamous cell carcinoma; chronic ulcerative stomatitis protein; keratinocyte transcription factor KET; tumor protein p63 deltaN isoform delta; AIS; LMS; RHS; p40; TP53L; TP73L; TP53CP; B(p51A); B(p51B);

Gene ID

8626

mRNA Refseq

NM_001114978

Protein Refseq

NP_001108450

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	603273
UniProt ID	Q9H3D4
Chromosome Location	3q27-q29
Pathway	Apoptosis, organism-specific biosystem; Direct p53 effectors, organism-specific biosystem; TP53 network, organism-specific biosystem;
Function	DNA binding; chromatin binding; chromatin binding; damaged DNA binding; double-stranded DNA binding; identical protein binding; metal ion binding; p53 binding; protein binding; sequence-specific DNA binding; sequence-specific DNA binding transcription factor activity; transcription regulatory region DNA binding;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA