

Recombinant Human NAMPT cell lysate

Cat. No. NAMPT-491HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human NAMPT / PBEF derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	NAMPT nicotinamide phosphoribosyltransferase [Homo sapiens]
Official Symbol	NAMPT
Synonyms	NAMPT; nicotinamide phosphoribosyltransferase; PBEF1, pre B cell colony enhancing factor 1; PBEF; visfatin; NAMPRTase; pre-B cell-enhancing factor; pre-B-cell colony enhancing factor 1; pre-B-cell colony-enhancing factor 1; VF; PBEF1; VISFATIN; 1110035O14Rik; MGC117256; DKFZp666B131;
Gene ID	10135
mRNA Refseq	NM_005746
Protein Refseq	NP_005737
MIM	608764
UniProt ID	P43490

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	7q22.2
Pathway	Adipogenesis, organism-specific biosystem; BMAL1:CLOCK/NPAS2 Activates Circadian Expression, organism-specific biosystem; Circadian Clock, organism-specific biosystem; Metabolism, organism-specific biosystem; Metabolism of vitamins and cofactors, organism-specific biosystem; Metabolism of water-soluble vitamins and cofactors, organism-specific biosystem; NAD biosynthesis III, organism-specific biosystem;
Function	cytokine activity; nicotinamide phosphoribosyltransferase activity; nicotinate phosphoribosyltransferase activity; nicotinate-nucleotide diphosphorylase (carboxylating) activity; transferase activity, transferring glycosyl groups;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA