

Recombinant Human PPARG cell lysate

Cat. No. PPARG-492HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human PPARG / Nr1c3 / PPARGgamma derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.
GENE INFORMATION	
Gene Name	PPARG peroxisome proliferator-activated receptor gamma [Homo sapiens]
Official Symbol	PPARG
Synonyms	PPARG; peroxisome proliferator-activated receptor gamma; peroxisome proliferative activated receptor, gamma; NR1C3; PPARG1; PPARG2; PPARGgamma; PPAR gamma; PPAR-gamma; nuclear receptor subfamily 1 group C member 3; peroxisome proliferator-activated receptor gamma 1; peroxisome proliferator-activated nuclear receptor gamma variant 1; GLM1; CIMT1;
Gene ID	5468
mRNA Refseq	NM_005037
Protein Refseq	NP_005028
MIM	601487

UniProt ID	P37231
Chromosome Location	3p25
Pathway	Adipogenesis, organism-specific biosystem; Calcineurin-regulated NFAT-dependent transcription in lymphocytes, organism-specific biosystem; Developmental Biology, organism-specific biosystem; Energy Metabolism, organism-specific biosystem; Fatty acid, triacylglycerol, and ketone body metabolism, organism-specific biosystem; Gene Expression, organism-specific biosystem; Generic Transcription Pathway, organism-specific biosystem;
Function	DNA binding; RNA polymerase II regulatory region DNA binding; activating transcription factor binding; arachidonic acid binding; chromatin binding; drug binding; enzyme binding; ligand-activated sequence-specific DNA binding RNA polymerase II transcription factor activity; ligand-activated sequence-specific DNA binding RNA polymerase II transcription factor activity; metal ion binding; prostaglandin receptor activity; protein binding; receptor activity; retinoid X receptor binding; contributes_to sequence-specific DNA binding; sequence-specific DNA binding; sequence-specific DNA binding transcription factor activity; sequence-specific DNA binding transcription factor activity; sequence-specific DNA binding transcription factor activity; steroid hormone receptor activity; transcription regulatory region DNA binding; transcription regulatory region DNA binding; zinc ion binding;