

Recombinant Human ATF2 cell lysate

Cat. No. ATF2-518HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human ATF2 derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

[ATF2 activating transcription factor 2 \[Homo sapiens \]](#)

Official Symbol

ATF2

Synonyms

ATF2; activating transcription factor 2; cAMP responsive element binding protein 2 , CREB2; cyclic AMP-dependent transcription factor ATF-2; CRE BP1; HB16; TREB7; CREB-2; cAMP-dependent transcription factor ATF-2; cAMP-responsive element-binding protein 2; cAMP response element-binding protein CRE-BP1; cyclic AMP-responsive element-binding protein 2; cAMP responsive element binding protein 2, formerly; activating transcription factor 2 splice variant ATF2-var2; CREB2; CRE-BP1; MGC111558;

Gene ID

[1386](#)

mRNA Refseq

[NM_001256090](#)

Protein Refseq

[NP_001243019](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	123811
UniProt ID	P15336
Chromosome Location	2q32
Pathway	ATF-2 transcription factor network, organism-specific biosystem; Activated TLR4 signalling, organism-specific biosystem; Activation of the AP-1 family of transcription factors, organism-specific biosystem; Amphetamine addiction, organism-specific biosystem; Amphetamine addiction, conserved biosystem; Androgen Receptor Signaling Pathway, organism-specific biosystem; B Cell Receptor Signaling Pathway, organism-specific biosystem;
Function	DNA binding; RNA polymerase II activating transcription factor binding; RNA polymerase II core promoter proximal region sequence-specific DNA binding transcription factor activity involved in positive regulation of transcription; RNA polymerase II distal enhancer sequence-specific DNA binding; RNA polymerase II transcription factor binding transcription factor activity; cAMP response element binding; cAMP response element binding protein binding; chromatin binding; metal ion binding; protein binding; protein dimerization activity; protein kinase binding; sequence-specific DNA binding; sequence-specific DNA binding transcription factor activity; sequence-specific DNA binding transcription factor activity; sequence-specific distal enhancer binding RNA polymerase II transcription factor activity; transcription coactivator activity; zinc ion binding;