

Recombinant Rat VEGFA cell lysate

Cat. No. VEGFA-522RCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Rat VEGF164 / VEGFA derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Rat
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	Vegfa vascular endothelial growth factor A [Rattus norvegicus]
Official Symbol	VEGFA
Synonyms	VEGFA; vascular endothelial growth factor A; VPF; VEGF-A; vascular permeability factor; Vegf; VEGF164;
Gene ID	83785
mRNA Refseq	NM_001110333
Protein Refseq	NP_001103803
Pathway	Bladder cancer, organism-specific biosystem; Bladder cancer, conserved biosystem; Endochondral Ossification, organism-specific biosystem; Focal adhesion, organism-specific biosystem; Focal adhesion, conserved biosystem; Hemostasis, organism-specific biosystem; Hypertrophy Model, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Function

cell surface binding; cell surface binding; chemoattractant activity; chemoattractant activity; cytokine activity; cytokine activity; fibronectin binding; fibronectin binding; growth factor activity; growth factor activity; heparin binding; heparin binding; platelet-derived growth factor receptor binding; platelet-derived growth factor receptor binding; protein heterodimerization activity; protein heterodimerization activity; protein homodimerization activity; receptor agonist activity; vascular endothelial growth factor receptor 1 binding; vascular endothelial growth factor receptor 1 binding; vascular endothelial growth factor receptor 2 binding; vascular endothelial growth factor receptor 2 binding; vascular endothelial growth factor receptor 2 binding; vascular endothelial growth factor receptor binding; vascular endothelial growth factor receptor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA