

Recombinant Mouse PLK1 cell lysate

Cat. No. PLK1-532MCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Mouse PLK1 / PLK-1 derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Mouse
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	Plk1 polo-like kinase 1 (Drosophila) [Mus musculus]
Official Symbol	PLK1
Synonyms	PLK1; polo-like kinase 1 (Drosophila); serine/threonine-protein kinase PLK1; PLK-1; serine/threonine-protein kinase 13; Plk; STPK13;
Gene ID	18817
mRNA Refseq	NM_011121
Protein Refseq	NP_035251
Pathway	APC/C-mediated degradation of cell cycle proteins, organism-specific biosystem; APC/C:Cdh1 mediated degradation of Cdc20 and other APC/C:Cdh1 targeted proteins in late mitosis/early G1, organism-specific biosystem; Activation of APC/C and APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem; Cell Cycle, organism-specific biosystem; Cell Cycle, Mitotic, organism-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

specific biosystem; Cell cycle, organism-specific biosystem; Cell cycle, organism-specific biosystem;

Function

ATP binding; anaphase-promoting complex binding; kinase activity; microtubule binding; nucleotide binding; protein kinase activity; protein kinase binding; protein serine/threonine kinase activity; transferase activity; transferase activity, transferring phosphorus-containing groups;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA