

Recombinant Mouse IL5 protein

Cat. No. IL5-206M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Mouse IL5 protein was expressed in Escherichia coli.
Species	Mouse
Source	E.coli
ProteinLength	113
Description	This gene encodes a cytokine that acts as a growth and differentiation factor for both B cells and eosinophils. The encoded cytokine plays a major role in the regulation of eosinophil formation, maturation, recruitment and survival. The increased production of this cytokine may be related to pathogenesis of eosinophil-dependent inflammatory diseases. This cytokine functions by binding to its receptor, which is a heterodimer, whose beta subunit is shared with the receptors for interleukine 3 (IL3) and colony stimulating factor 2 (CSF2/GM-CSF). This gene is located on chromosome 5 within a cytokine gene cluster which includes interleukin 4 (IL4), interleukin 13 (IL13), and CSF2 .
Form	Lyophilized from a 0.2µm filtered concentrated solution in 20 mM Tris, pH 9.0, 150 mM NaCl.
Bio-activity	Fully biologically active when compared to standard. The ED50 as determined by a cell proliferation assay using human TF-1 cells is less than 2 ng/ml, corresponding to a specific activity of $> 5.0 \times 10^5$ IU/mg.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	Approximately 26.2 kDa, a disulfide-linked homodimeric protein containing two 113 amino acids.
AA Sequence	MEIPMSTVVKETLTQLSAHRALLTSNETMRLPVPTHKNHQLCIGEIFQGLDILKNQTV RGGTVEMLFQNLSLIKKYIDRQKEKCGEERRRTRQFLDYQLQEFLGVMSTEWAMEG
Endotoxin	Less than 0.1 EU/μg of rMuLL-5 as determined by LAL method.
Purity	>98% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	IL5
Official Symbol	IL5
Synonyms	IL5; interleukin 5; interleukin-5; TRF; BCGF-II; B-cell growth factor II; T-cell replacing factor; cytotoxic T-lymphocyte inducer; eosinophil differentiation factor; IL-5;
Gene ID	16191

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

mRNA Refseq NM_010558

Protein Refseq NP_034688

UniProt ID P04401

SDS-PAGE of IL5-
206M



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA