

Recombinant Mouse PRKCD cell lysate

Cat. No. PRKCD-546MCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Mouse PKC delta / PRKCD derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Mouse
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	Prkcd protein kinase C, delta [Mus musculus]
Official Symbol	PRKCD
Synonyms	PRKCD; protein kinase C, delta; protein kinase C delta type; nPKC-delta; protein kinase C[d]; protein kinase C, delta V; protein kinase C, delta IV; tyrosine-protein kinase PRKCD; protein kinase C delta variant IX; Pkcd; PKC[d]; AI385711; PKCdelta; D14Ert420e;
Gene ID	18753
mRNA Refseq	NM_011103
Protein Refseq	NP_035233
Pathway	Alpha6-Beta4 Integrin Signaling Pathway, organism-specific biosystem; Apoptosis, organism-specific biosystem; Apoptotic cleavage of cellular proteins, organism-specific biosystem; Apoptotic executionphase, organism-specific biosystem; B Cell

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Receptor Signaling Pathway, organism-specific biosystem; Ca-dependent events, organism-specific biosystem; CaM pathway, organism-specific biosystem;

Function

ATP binding; TIR domain binding; calcium-independent protein kinase C activity; enzyme activator activity; enzyme binding; insulin receptor substrate binding; kinase activity; metal ion binding; non-membrane spanning protein tyrosine kinase activity; nucleotide binding; protein C-terminus binding; protein kinase C activity; protein kinase activity; protein kinase binding; protein serine/threonine kinase activity; transferase activity; transferase activity, transferring phosphorus-containing groups;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA