

Recombinant Human STAT1 cell lysate

Cat. No. STAT1-555HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human STAT1 / p91 derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

[STAT1 signal transducer and activator of transcription 1, 91kDa \[Homo sapiens \]](#)

Official Symbol

STAT1

Synonyms

STAT1; signal transducer and activator of transcription 1, 91kDa; signal transducer and activator of transcription 1, 91kD; signal transducer and activator of transcription 1-alpha/beta; ISGF 3; STAT91; transcription factor ISGF 3 components p91/p84; transcription factor ISGF-3 components p91/p84; signal transducer and activator of transcription-1; CANDF7; ISGF-3; DKFZp686B04100;

Gene ID

[6772](#)

mRNA Refseq

[NM_007315](#)

Protein Refseq

[NP_009330](#)

MIM

[600555](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	P42224
Chromosome Location	2q32.2-q32.3
Pathway	Adipogenesis, organism-specific biosystem; Antiviral mechanism by IFN-stimulated genes, organism-specific biosystem; B Cell Receptor Signaling Pathway, organism-specific biosystem; CXCR4-mediated signaling events, organism-specific biosystem; Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Cytokine Signaling in Immune system, organism-specific biosystem;
Function	RNA polymerase II core promoter sequence-specific DNA binding; RNA polymerase II core promoter sequence-specific DNA binding transcription factor activity; calcium ion binding; double-stranded DNA binding; enzyme binding; non-membrane spanning protein tyrosine kinase activity; protein binding; protein homodimerization activity; sequence-specific DNA binding; sequence-specific DNA binding transcription factor activity; NOT sequence-specific DNA binding transcription factor activity; signal transducer activity; tumor necrosis factor receptor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA