

Recombinant Human RHOA cell lysate

Cat. No. RHOA-601HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human RHOA derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

RHOA ras homolog family member A [*Homo sapiens*]

Official Symbol

RHOA

Synonyms

RHOA; ras homolog family member A; ARH12, ARHA, ras homolog gene family, member A; transforming protein RhoA; Rho12; RhoA; RHOH12; h12; oncogene RHO H12; rho cDNA clone 12; Aplysia ras-related homolog 12; small GTP binding protein RhoA; ras homolog gene family, member A; ARHA; ARH12; RHO12;

Gene ID

387

mRNA Refseq

NM_001664

Protein Refseq

NP_001655

MIM

165390

UniProt ID

P61586

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	3p21.3
Pathway	Adherens junction, organism-specific biosystem; Adherens junction, conserved biosystem; Alpha6-Beta4 Integrin Signaling Pathway, organism-specific biosystem; Arf6 downstream pathway, organism-specific biosystem; Aurora B signaling, organism-specific biosystem; Axon guidance, organism-specific biosystem; Axon guidance, conserved biosystem;
Function	GTP binding; GTPase activity; myosin binding; nucleotide binding; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA