

Recombinant Human PDK1 cell lysate

Cat. No. PDK1-603HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human PDK-1 derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	PDK1 pyruvate dehydrogenase kinase, isozyme 1 [Homo sapiens]
Official Symbol	PDK1
Synonyms	PDK1; pyruvate dehydrogenase kinase, isozyme 1; pyruvate dehydrogenase kinase, isoenzyme 1; mitochondrial pyruvate dehydrogenase, lipoamide, kinase isoenzyme 1;
Gene ID	5163
mRNA Refseq	NM_002610
Protein Refseq	NP_002601
MIM	602524
UniProt ID	Q15118
Chromosome Location	2q31.1

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway

EPO Receptor Signaling, organism-specific biosystem; ErbB signaling pathway, organism-specific biosystem; Fc epsilon RI signaling pathway, organism-specific biosystem; Fc epsilon RI signaling pathway, conserved biosystem; Hepatitis C, organism-specific biosystem; Hepatitis C, conserved biosystem; Metabolism, organism-specific biosystem;

Function

ATP binding; nucleotide binding; protein complex binding; protein heterodimerization activity; protein homodimerization activity; protein kinase activity; pyruvate dehydrogenase (acetyl-transferring) kinase activity; transferase activity; two-component sensor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA