

Recombinant Human FES cell lysate

Cat. No. FES-606HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human FES Kinase / Feline sarcoma oncogene derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.
GENE INFORMATION	
Gene Name	FES feline sarcoma oncogene [Homo sapiens]
Official Symbol	FES
Synonyms	FES; feline sarcoma oncogene; feline sarcoma (Snyder Theilen) viral (v fes)/Fujinami avian sarcoma (PRCII) viral (v fps) oncogene homolog; tyrosine-protein kinase Fes/Fps; c fes/fps protein; FPS; Oncogene FES; feline sarcoma virus; p93c-fes; proto-oncogene c-Fes; proto-oncogene c-Fps; Oncogene FES, feline sarcoma virus; proto-oncogene tyrosine-protein kinase Fes/Fps; feline sarcoma/Fujinami avian sarcoma oncogene homolog; feline sarcoma (Snyder-Theilen) viral (v-fes)/Fujinami avian sarcoma (PRCII) viral (v-fps) oncogene homolog;
Gene ID	2242
mRNA Refseq	NM_001143783
Protein Refseq	NP_001137255

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	190030
UniProt ID	P07332
Chromosome Location	15q25-qter
Pathway	Angiopoietin receptor Tie2-mediated signaling, organism-specific biosystem; Axon guidance, organism-specific biosystem; Axon guidance, conserved biosystem; Axon guidance, organism-specific biosystem; CRMPs in Sema3A signaling, organism-specific biosystem; Developmental Biology, organism-specific biosystem; IL-3 Signaling Pathway, organism-specific biosystem;
Function	ATP binding; immunoglobulin receptor binding; lipid binding; non-membrane spanning protein tyrosine kinase activity; nucleotide binding; phosphatidylinositol binding; protein tyrosine kinase activity;