

Recombinant Human MAP2K2 cell lysate

Cat. No. MAP2K2-613HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human MAP2K2 / MEK2 derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

MAP2K2 mitogen-activated protein kinase kinase 2 [Homo sapiens]

Official Symbol

MAP2K2

Synonyms

MAP2K2; mitogen-activated protein kinase kinase 2; PRKMK2; dual specificity mitogen-activated protein kinase kinase 2; MEK2; MAPK/ERK kinase 2; MAP kinase kinase 2; ERK activator kinase 2; mitogen-activated protein kinase kinase 2, p45; MKK2; MAPKK2; FLJ26075;

Gene ID

5605

mRNA Refseq

NM_030662

Protein Refseq

NP_109587

MIM

601263

UniProt ID

P36507

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	19p13.3
Pathway	ARMS-mediated activation, organism-specific biosystem; Activated TLR4 signalling, organism-specific biosystem; Acute myeloid leukemia, organism-specific biosystem; Acute myeloid leukemia, conserved biosystem; Axon guidance, organism-specific biosystem; B Cell Receptor Signaling Pathway, organism-specific biosystem; B cell receptor signaling pathway, organism-specific biosystem;
Function	ATP binding; nucleotide binding; protein binding; protein complex scaffold; protein serine/threonine kinase activity; protein serine/threonine kinase activity; protein serine/threonine/tyrosine kinase activity; protein tyrosine kinase activity;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA