

Recombinant Human PDE4B cell lysate

Cat. No. PDE4B-621HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human PDE4B / DPDE4 derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

PDE4B phosphodiesterase 4B, cAMP-specific [Homo sapiens]

Official Symbol

PDE4B

Synonyms

PDE4B; phosphodiesterase 4B, cAMP-specific; DPDE4, phosphodiesterase 4B, cAMP specific (dunce (Drosophila) homolog phosphodiesterase E4) , phosphodiesterase 4B, cAMP specific (phosphodiesterase E4 dunce homolog, Drosophila); cAMP-specific 3,5-cyclic phosphodiesterase 4B; phosphodiesterase E4 dunce homolog (Drosophila); PDE32; dunce-like phosphodiesterase E4; cAMP-specific phosphodiesterase-4 B isoform; phosphodiesterase 4B, cAMP-specific (phosphodiesterase E4 dunce homolog, Drosophila); DPDE4; PDE4B5; PDEIVB; MGC126529; DKFZp686F2182;

Gene ID

5142

mRNA Refseq

NM_001037339

Protein Refseq

NP_001032416

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	600127
UniProt ID	Q07343
Chromosome Location	1p31
Pathway	DARPP-32 events, organism-specific biosystem; G Protein Signaling Pathways, organism-specific biosystem; G alpha (s) signalling events, organism-specific biosystem; GPCR downstream signaling, organism-specific biosystem; Morphine addiction, organism-specific biosystem; Morphine addiction, conserved biosystem; Myometrial Relaxation and Contraction Pathways, organism-specific biosystem;
Function	3,5-cyclic-AMP phosphodiesterase activity; hydrolase activity; metal ion binding; phosphoric diester hydrolase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA