

Recombinant Human MAP4K5 cell lysate

Cat. No. MAP4K5-631HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human MAP4K5 / MEKKK5 derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	MAP4K5 mitogen-activated protein kinase kinase kinase kinase 5 [Homo sapiens]
Official Symbol	MAP4K5
Synonyms	MAP4K5; mitogen-activated protein kinase kinase kinase kinase 5; GCKR; germinal center kinase related; KHS; KHS1; MEKKK 5; MEK kinase kinase 5; germinal center kinase-related; MAPK/ERK kinase kinase kinase 5; kinase homologous to SPS1/STE20; MAPKKKK5;
Gene ID	11183
mRNA Refseq	NM_006575
Protein Refseq	NP_006566
MIM	604923
UniProt ID	Q9Y4K4

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	14q11.2-q21
Pathway	Insulin Signaling, organism-specific biosystem; TNF receptor signaling pathway, organism-specific biosystem;
Function	ATP binding; nucleotide binding; protein binding; protein kinase activity; protein serine/threonine kinase activity; small GTPase regulator activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA