

Recombinant Mouse CNDP2 cell lysate

Cat. No. CNDP2-641MCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Mouse CNDP2 / CN2 / CPGL derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Mouse
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.
GENE INFORMATION	
Gene Name	Cndp2 CNDP dipeptidase 2 (metallopeptidase M20 family) [<i>Mus musculus</i>]
Official Symbol	CNDP2
Synonyms	CNDP2; CNDP dipeptidase 2 (metallopeptidase M20 family); cytosolic non-specific dipeptidase; peptidase 1; peptidase A; cytosolic nonspecific dipeptidase; glutamate carboxypeptidase-like protein 1; Cn2; Pep1; Dip-2; Pep-1; C76600; 0610010E05Rik;
Gene ID	66054
mRNA Refseq	NM_023149
Protein Refseq	NP_075638
Pathway	Arginine and proline metabolism, organism-specific biosystem; Arginine and proline metabolism, conserved biosystem; Histidine metabolism, organism-specific biosystem; Histidine metabolism, conserved biosystem; Metabolic pathways,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

organism-specific biosystem; beta-Alanine metabolism, organism-specific biosystem;
beta-Alanine metabolism, conserved biosystem;

Function

carboxypeptidase activity; dipeptidase activity; dipeptidase activity; hydrolase activity;
metal ion binding; metallopeptidase activity; peptidase activity; tripeptidase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA