

Recombinant Human USP7 cell lysate

Cat. No. USP7-671HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human USP7 / HAUSP derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

USP7 ubiquitin specific peptidase 7 (herpes virus-associated) [Homo sapiens]

Official Symbol

USP7

Synonyms

USP7; ubiquitin specific peptidase 7 (herpes virus-associated); HAUSP, ubiquitin specific protease 7 (herpes virus associated); ubiquitin carboxyl-terminal hydrolase 7; ubiquitin thioesterase 7; deubiquitinating enzyme 7; ubiquitin-specific-processing protease 7; herpesvirus-associated ubiquitin-specific protease; Herpes virus-associated ubiquitin-specific protease; ubiquitin specific protease 7 (herpes virus-associated); TEF1; HAUSP;

Gene ID

7874

mRNA Refseq

NM_003470

Protein Refseq

NP_003461

MIM

602519

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	Q93009
Chromosome Location	16p13.3
Pathway	FoxO family signaling, organism-specific biosystem; Herpes simplex infection, organism-specific biosystem; Herpes simplex infection, conserved biosystem; p53 pathway, organism-specific biosystem;
Function	cysteine-type endopeptidase activity; p53 binding; peptidase activity; protein C-terminus binding; protein binding; protein homodimerization activity; transcription factor binding; ubiquitin protein ligase binding; ubiquitin thiolesterase activity; ubiquitin-specific protease activity; ubiquitin-specific protease activity;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA