

## Recombinant Human TGM2 cell lysate

**Cat. No.** TGM2-686HCL    **Lot. No.** (See product label)

### SPECIFICATION

|                                |                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|--------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Product Overview</b>        | Human TGM2 / TGC / Transglutaminase-2 derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls                                                                                                                                                                                                                                                              |
| <b>Species</b>                 | Human                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| <b>Source</b>                  | Insect Cells                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| <b>Preparation method</b>      | Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized. |
| <b>Lysis buffer</b>            | Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF                                                                                                                                                                                                                                                                                                               |
| <b>Quality control Testing</b> | 12.5% SDS-PAGE Stained with Coomassie Blue                                                                                                                                                                                                                                                                                                                                                                                                          |
| <b>Recommended Usage</b>       | 1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on                                                                                                 |

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

|                            |                                                                                                                                                                                                                                                                                                                                                              |
|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                            | SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.                                                                                                                                                                                                                                         |
| <b>Stability</b>           | Samples are stable for up to twelve months from date of receipt at -80°C                                                                                                                                                                                                                                                                                     |
| <b>Storage Buffer</b>      | 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF                                                                                                                                                                                                                                                    |
| <b>Storage Instruction</b> | Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.                                                                                                                                                                        |
| <b>GENE INFORMATION</b>    |                                                                                                                                                                                                                                                                                                                                                              |
| <b>Gene Name</b>           | TGM2 transglutaminase 2 (C polypeptide, protein-glutamine-gamma-glutamyltransferase) [ Homo sapiens ]                                                                                                                                                                                                                                                        |
| <b>Official Symbol</b>     | TGM2                                                                                                                                                                                                                                                                                                                                                         |
| <b>Synonyms</b>            | TGM2; transglutaminase 2 (C polypeptide, protein-glutamine-gamma-glutamyltransferase); protein-glutamine gamma-glutamyltransferase 2; TGC; TG(C); TGase C; TGase H; TGase-2; TGase-H; C polypeptide; transglutaminase C; transglutaminase H; transglutaminase-2; tissue transglutaminase; protein-glutamine-gamma-glutamyltransferase; TG2; GNAH; G-ALPHA-h; |
| <b>Gene ID</b>             | 7052                                                                                                                                                                                                                                                                                                                                                         |
| <b>mRNA Refseq</b>         | NM_004613                                                                                                                                                                                                                                                                                                                                                    |
| <b>Protein Refseq</b>      | NP_004604                                                                                                                                                                                                                                                                                                                                                    |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

|                            |                                                                                                                                                                                         |
|----------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>MIM</b>                 | 190196                                                                                                                                                                                  |
| <b>UniProt ID</b>          | P21980                                                                                                                                                                                  |
| <b>Chromosome Location</b> | 20q12                                                                                                                                                                                   |
| <b>Pathway</b>             | Huntingtons disease, organism-specific biosystem; Huntingtons disease, conserved biosystem; Thromboxane A2 receptor signaling, organism-specific biosystem;                             |
| <b>Function</b>            | GTP binding; metal ion binding; protein binding; protein domain specific binding; protein-glutamine gamma-glutamyltransferase activity; transferase activity, transferring acyl groups; |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA