

Recombinant Human S100A6 cell lysate

Cat. No. S100A6-690HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human S100A6 / CACY derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

[S100A6](#) [S100 calcium binding protein A6 \[Homo sapiens \]](#)

Official Symbol

[S100A6](#)

Synonyms

[S100A6](#); [S100 calcium binding protein A6](#); [CACY](#), [S100 calcium binding protein A6 \(calcyclin\)](#) , [S100 calcium binding protein A6 \(calcyclin\)](#); [protein S100-A6](#); [2A9](#); [CABP](#); [PRA](#); [MLN 4](#); [calcyclin](#); [growth factor-inducible protein 2A9](#); [prolactin receptor-associated protein](#); [S100 calcium-binding protein A6 \(calcyclin\)](#); [5B10](#); [CACY](#);

Gene ID

[6277](#)

mRNA Refseq

[NM_014624](#)

Protein Refseq

[NP_055439](#)

MIM

[114110](#)

UniProt ID

[P06703](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	1q21
Pathway	Prostaglandin Synthesis and Regulation, organism-specific biosystem;
Function	S100 beta binding; calcium ion binding; calcium ion binding; calcium-dependent protein binding; ion transmembrane transporter activity; protein binding; protein homodimerization activity; protein homodimerization activity; tropomyosin binding; zinc ion binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA