

Recombinant Human MAPK1 cell lysate

Cat. No. MAPK1-692HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human MAPK1 / ERK2 derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

[MAPK1 mitogen-activated protein kinase 1 \[Homo sapiens \]](#)

Official Symbol

MAPK1

Synonyms

MAPK1; mitogen-activated protein kinase 1; PRKM1, PRKM2; ERK; ERK2; MAPK2; p41mapk; ERK-2; MAPK 2; p42-MAPK; MAP kinase 1; MAP kinase 2; MAP kinase isoform p42; protein tyrosine kinase ERK2; mitogen-activated protein kinase 2; extracellular signal-regulated kinase 2; p38; p40; p41; ERT1; PRKM1; PRKM2; P42MAPK;

Gene ID

[5594](#)

mRNA Refseq

[NM_002745](#)

Protein Refseq

[NP_002736](#)

MIM

[176948](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	P28482
Chromosome Location	22q11.2
Pathway	ALK1 signaling events, organism-specific biosystem; ARMS-mediated activation, organism-specific biosystem; ATF-2 transcription factor network, organism-specific biosystem; Activated TLR4 signalling, organism-specific biosystem; Activation of NMDA receptor upon glutamate binding and postsynaptic events, organism-specific biosystem; Activation of the AP-1 family of transcription factors, organism-specific biosystem; Acute myeloid leukemia, organism-specific biosystem;
Function	ATP binding; DNA binding; MAP kinase activity; RNA polymerase II carboxy-terminal domain kinase activity; mitogen-activated protein kinase kinase kinase binding; nucleotide binding; phosphatase binding; phosphotyrosine binding; protein binding; protein serine/threonine kinase activity; transcription factor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA