

Recombinant Human HEXB cell lysate

Cat. No. HEXB-756HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human HEXB / Hexosaminidase B derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

HEXB hexosaminidase B (beta polypeptide) [Homo sapiens]

Official Symbol

HEXB

Synonyms

HEXB; hexosaminidase B (beta polypeptide); beta-hexosaminidase subunit beta; HCC-7; hexosaminidase subunit B; beta-N-acetylhexosaminidase subunit beta; cervical cancer proto-oncogene 7 protein; N-acetyl-beta-glucosaminidase subunit beta; ENC-1AS;

Gene ID

3074

mRNA Refseq

NM_000521

Protein Refseq

NP_000512

MIM

606873

UniProt ID

P07686

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	5q13.3
Pathway	Amino sugar and nucleotide sugar metabolism, organism-specific biosystem; Amino sugar and nucleotide sugar metabolism, conserved biosystem; Glycosaminoglycan degradation, organism-specific biosystem; Glycosaminoglycan degradation, conserved biosystem; Glycosphingolipid biosynthesis - ganglio series, organism-specific biosystem; Glycosphingolipid biosynthesis - ganglio series, conserved biosystem; Glycosphingolipid biosynthesis - globo series, organism-specific biosystem;
Function	beta-N-acetylglucosaminidase activity; beta-N-acetylhexosaminidase activity; catalytic activity; cation binding; protein heterodimerization activity; protein heterodimerization activity; protein homodimerization activity; protein homodimerization activity; sugar binding;