

Recombinant Human CD47 cell lysate

Cat. No. CD47-850HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human CD47 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	CD47 CD47 molecule [Homo sapiens]
Official Symbol	CD47
Synonyms	CD47; CD47 molecule; CD47 antigen (Rh related antigen, integrin associated signal transducer) , MER6; leukocyte surface antigen CD47; antigen identified by monoclonal 1D8; antigenic surface determinant protein OA3; CD47 glycoprotein; IAP; integrin associated protein; OA3; Rh related antigen; Rh-related antigen; integrin-associated protein; integrin-associated signal transducer; CD47 antigen (Rh-related antigen, integrin-associated signal transducer); MER6;
Gene ID	961
mRNA Refseq	NM_001777
Protein Refseq	NP_001768
MIM	601028

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	Q08722
Chromosome Location	3q13.1-q13.2
Pathway	Cell surface interactions at the vascular wall, organism-specific biosystem; Cell-Cell communication, organism-specific biosystem; ECM-receptor interaction, organism-specific biosystem; ECM-receptor interaction, conserved biosystem; Hemostasis, organism-specific biosystem; Integrins in angiogenesis, organism-specific biosystem; Signal regulatory protein (SIRP) family interactions, organism-specific biosystem;
Function	protein binding; thrombospondin receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA