

Recombinant Human EDEM2 cell lysate

Cat. No. EDEM2-862HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human EDEM2 / C20orf31 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	EDEM2 ER degradation enhancer, mannosidase alpha-like 2 [Homo sapiens]
Official Symbol	EDEM2
Synonyms	EDEM2; ER degradation enhancer, mannosidase alpha-like 2; C20orf31, C20orf49, chromosome 20 open reading frame 31; ER degradation-enhancing alpha-mannosidase-like 2; bA4204.1; FLJ10783; ER degradation-enhancing-mannosidase-like protein 2; C20orf31; C20orf49;
Gene ID	55741
mRNA Refseq	NM_001145025
Protein Refseq	NP_001138497
MIM	610302
UniProt ID	Q9BV94

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	20q11.22
Pathway	Asparagine N-linked glycosylation, organism-specific biosystem; Calnexin/calreticulin cycle, organism-specific biosystem; ER Quality Control Compartment (ERQC), organism-specific biosystem; Metabolism of proteins, organism-specific biosystem; N-glycan trimming in the ER and Calnexin/Calreticulin cycle, organism-specific biosystem; Post-translational protein modification, organism-specific biosystem; Protein processing in endoplasmic reticulum, organism-specific biosystem;
Function	calcium ion binding; NOT mannosyl-oligosaccharide 1,2-alpha-mannosidase activity;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA