

Recombinant Human LILRA3 cell lysate

Cat. No. LILRA3-976HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human LILRA3 / ILT6 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	LILRA3 leukocyte immunoglobulin-like receptor, subfamily A (without TM domain), member 3 [Homo sapiens]
Official Symbol	LILRA3
Synonyms	LILRA3; leukocyte immunoglobulin-like receptor, subfamily A (without TM domain), member 3; leukocyte immunoglobulin-like receptor subfamily A member 3; CD85e; HM31; HM43; ILT6; LIR 4; LIR4; ILT-6; immunoglobulin-like transcript 6; CD85 antigen-like family member E; leucocyte immunoglobulin-like receptor; monocyte inhibitory receptor HM43/HM31; leukocyte immunoglobulin-like receptor 4; leukocyte immunoglobulin-like receptor A3; e3; CD85E; LIR-4;
Gene ID	11026
mRNA Refseq	NM_001172654
Protein Refseq	NP_001166125

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	604818
UniProt ID	Q8N6C8
Chromosome Location	19q13.4
Pathway	Osteoclast differentiation, organism-specific biosystem; Osteoclast differentiation, conserved biosystem;
Function	antigen binding; receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA