

Recombinant Mouse CD3D & CD3E cell lysate

Cat. No. CD3D & CD3E-1076MCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Mouse CD3D & CD3E Heterodimer derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Mouse
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	Cd3d CD3 antigen, delta polypeptide [<i>Mus musculus</i>]; Cd3e CD3 antigen, epsilon polypeptide [<i>Mus musculus</i>]
Official Symbol	Cd3d; Cd3e
Synonyms	T3d; T-cell surface glycoprotein CD3 delta chain; T-cell receptor T3 delta chain; CD3; T3e; AI504783; CD3epsilon; T-cell surface glycoprotein CD3 epsilon chain; T-cell surface antigen T3/Leu-4 epsilon chain
Gene ID	12500; 12501
mRNA Refseq	NM_013487; NM_007648
Protein Refseq	NP_038515; NP_031674
UniProt ID	P04235; P22646
Chromosome	9 A5.2; 9 24.84 cM; 9 A5.2; 9 24.84 cM

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Location**Pathway**

Adaptive Immune System, organism-specific biosystem; Hematopoietic cell lineage, organism-specific biosystem; PD-1 signaling, organism-specific biosystem. Adaptive Immune System, organism-specific biosystem; Generation of second messenger molecules, organism-specific biosystem; Hematopoietic cell lineage, organism-specific biosystem

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA