

Recombinant Human SIGIRR cell lysate

Cat. No. SIGIRR-1091HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human SIGIRR / TIR8 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	SIGIRR single immunoglobulin and toll-interleukin 1 receptor (TIR) domain [Homo sapiens]
Official Symbol	SIGIRR
Synonyms	SIGIRR; single immunoglobulin and toll-interleukin 1 receptor (TIR) domain; single Ig IL-1-related receptor; single immunoglobulin domain IL1R1 related; TIR8; toll/interleukin-1 receptor 8; single Ig IL-1R-related molecule; single immunoglobulin domain-containing IL1R-related protein; MGC110992;
Gene ID	59307
mRNA Refseq	NM_001135053
Protein Refseq	NP_001128525
MIM	605478

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	Q6IA17
Chromosome Location	11p15.5
Pathway	Activated TLR4 signalling, organism-specific biosystem; Immune System, organism-specific biosystem; Innate Immune System, organism-specific biosystem; MyD88:Mal cascade initiated on plasma membrane, organism-specific biosystem; Toll Like Receptor 2 (TLR2) Cascade, organism-specific biosystem; Toll Like Receptor 4 (TLR4) Cascade, organism-specific biosystem; Toll Like Receptor TLR1:TLR2 Cascade, organism-specific biosystem;
Function	protein binding; transmembrane signaling receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA