

Recombinant Human LILRB2 cell lysate

Cat. No. LILRB2-1211HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human LILRB2 / ILT4 / LIR-2 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.
GENE INFORMATION	
Gene Name	LILRB2 leukocyte immunoglobulin-like receptor, subfamily B (with TM and ITIM domains), member 2 [Homo sapiens]
Official Symbol	LILRB2
Synonyms	LILRB2; leukocyte immunoglobulin-like receptor, subfamily B (with TM and ITIM domains), member 2; leukocyte immunoglobulin-like receptor subfamily B member 2; CD85d; ILT4; LIR 2; LIR2; MIR 10; MIR10; ILT-4; Ig-like transcript 4; immunoglobulin-like transcript 4; CD85 antigen-like family member D; leukocyte immunoglobulin-like receptor 2; monocyte/macrophage immunoglobulin-like receptor 10; leukocyte immunoglobulin-like receptor subfamily B member 2 soluble isoform; leukocyte immunoglobulin-like receptor, subfamily A (with TM domain), member 6; CD85D; LIR-2; LILRA6; MIR-10;
Gene ID	10288
mRNA Refseq	NM_001080978

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq	NP_001074447
MIM	604815
UniProt ID	Q8N423
Chromosome Location	19q13.4
Pathway	Adaptive Immune System, organism-specific biosystem; Immune System, organism-specific biosystem; Immunoregulatory interactions between a Lymphoid and a non-Lymphoid cell, organism-specific biosystem; Osteoclast differentiation, organism-specific biosystem; Osteoclast differentiation, conserved biosystem;
Function	MHC class I protein binding; MHC class I protein binding; MHC class Ib protein binding; cell adhesion molecule binding; eukaryotic cell surface binding; inhibitory MHC class I receptor activity; inhibitory MHC class I receptor activity; protein binding; protein phosphatase 1 binding; receptor activity;