

Recombinant Rat CD36 cell lysate

Cat. No. CD36-1236RCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Rat CD36 / GP3B derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Rat
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.
GENE INFORMATION	
Gene Name	Cd36 CD36 molecule (thrombospondin receptor) [<i>Rattus norvegicus</i>]
Official Symbol	CD36
Synonyms	CD36; CD36 molecule (thrombospondin receptor); platelet glycoprotein 4; GPIV; PAS-4; GPIIIB; PAS IV; glycoprotein IIb; fatty acid translocase; platelet glycoprotein IV; adipocyte membrane protein; fatty acid transport protein; collagen type I receptor thrombospondin receptor; CD36 antigen (collagen type I receptor, thrombospondin receptor); Fat; MGC91634; MGC108510;
Gene ID	29184
mRNA Refseq	NM_031561
Protein Refseq	NP_113749
Pathway	Adipocytokine signaling pathway, organism-specific biosystem; Adipocytokine signaling pathway, conserved biosystem; ECM-receptor interaction, organism-specific

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

biosystem; ECM-receptor interaction, conserved biosystem; Fat digestion and absorption, organism-specific biosystem; Fat digestion and absorption, conserved biosystem; Hematopoietic cell lineage, organism-specific biosystem;

Function

Gram-negative bacterial cell surface binding; Gram-positive bacterial cell surface binding; fatty acid binding; high-density lipoprotein particle binding; lipid binding; lipoprotein particle binding; lipoteichoic acid receptor activity; low-density lipoprotein particle binding; low-density lipoprotein receptor activity; protein binding; receptor activity; thrombospondin receptor activity; transforming growth factor beta binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA